

Electronic Supplementary Material

Comparison of concentration methods for detection of hepatitis A virus in water samples

Yuting Qiao^{1,3}, Zhiwei Sui², Guoliang Hu³, Huabin Cao³, Guoxiang Yang⁴, Yong Li⁴, Yongsong Lei⁴, Lihua Zhao¹, Quanjiao Chen^{1✉}

1. CAS Key Laboratory of Special Pathogens and Biosafety, Wuhan Institute of Virology, Chinese Academy of Sciences, Wuhan 430071, China

2. National Institute of Metrology, Beijing 100013, China

3. College of Animal Science and Technology, Institute of Animal Population Health, Jiangxi Agricultural University, Nanchang 330045, China

4. Hubei Wildlife Epidemic Focus and Disease Surveillance Center, Wuhan 430075, China

Supporting information to DOI: 10.1007/s12250-016-3786-5

Table S1. The primers used for hepatitis A virus (HAV) detection with conventional and real-time RT-PCR

Product	Primer name	Primer sequence (5'-3')
HAV247bp	HAV-F ^a	GTTTTGCTCCTCTTTATCATGCTATG
	HAV-R ^a	GGAAATGTCTCAGGTACTTTCTTTG
HAV	HAV-F2	TGCTATGGATGTTACTAC
	HAV-R2	ATCTTTCATGGTTGTTATAC
HCV	HCV-F	TCT GCG GAA CCG GTG AGT A
	HCV-R	TCA GGC AGT ACC ACA AGG C
AIV	AIV-F	TAT GAG AAG TGA AGT GGA A
	AIV-R	GTG TAT GTT GTG GAA TGG
HIV	HIV-F	TGT GTG CCC GTC TGT TGT GT
	HIV-R	GAG TCC TGC GTC GAG AGA GC

Note: ^a The primer HAV-F/R was used according to Brooks (2005).




Figure S1. Amplification curve (A and B) and standard curve (C) for HAV obtained by real-time RT-PCR.