Electronic Supplementary Material

A quantitative RT-PCR assay for rapid detection of Eurasianlineage H10 subtype influenza A virus

Hailiang Sun¹, Jian-Li Xue¹, Elizabeth Bailey¹, Yifei Xu¹, Guoliang Hu², John Baroch³, Yi Zhang⁴, Lanny Pace⁴, Thomas J. DeLiberto³, Xiu-Feng Wan^{1⊠}

1. Department of Basic Sciences, College of Veterinary Medicine, Mississippi State University, Mississippi 39762, USA

2. College of Animal Science and Technology, Jiangxi Agricultural University, Nanchang 330045, China 3. Animal and Plant Health Inspection Service of the US Department of Agriculture, Fort Collins 80526, USA

4. Mississippi Veterinary Research and Diagnostic Laboratory, College of Veterinary Medicine, Mississippi State University, Mississippi 39762, USA

Supporting information to DOI: 10.1007/s12250-016-3826-1



Figure S1. Specificity of quantitative RT-PCR for the detection of Eurasian-lineage H10 influenza A virus. Line A represents the amplification for the reassortant virus A/chicken/Jiangxi/JXA132717/2013 (HA, NA) × A/Puerto Rico/8/1934 (H10N8) amplification; Line B represents the amplifications of other influenza virus subtypes, including H1, H5, H7, H9, and North American-lineage H10N7 viruses, as well as other non-influenza avian viruses (Newcastle disease virus, infectious bursal disease virus, infectious bronchitis virus, and reovirus). The control reaction was conducted by using DEPC-treated water as template. The number of cycles (Ct) was calculated based on the maximum baseline-corrected normalized reporter (dRn) threshold of 0.27. Each assay was performed in triplicate.

Table S1. Viruses used to determine the specificity of a Eurasianlineage H10 specific quantitative RT-PCR method

Host	Virus name
Human	A/Puerto Rico/8/1934(H1N1)
	A/California/04/2009(H1N1)
	A/Brisbane/10/2005(H3N2)
	A/Perth/16/2009(H3N2)
Avian	A/mallard/Oregon/A00544081/2008(H1N1)
	A/mallard/Wisconsin/A00751454/2009(H1N1)
	A/mallard/Oregon/A00300758/2008(H2N3)
	A/mallard/Oregon/A00320372/2007(H2N3)
	A/mallard/Wisconsin/A00661712/2009(H3N2)
	A/mallard/Wisconsin/A00714818/2009(H3N2)
	A/mallard/Colorado/A00170379/2006(H3N8)
	A/mallard/Wisconsin/A00170366/2006(H3N8)
	A/mallard/Washington/A00714770/2009(H4N6)
	A/mallard/Washington/A00757090/2009(H4N6)
	A/chicken/Jiangxi/JXA132321/2013(H5N?) ^a
	A/mallard/Colorado/A00551322/2007(H6N1)
	A/mallard/New York/A00603000/2009(H6N1)
	A/blue-winged teal/Illinois/10OS3610/2010(H6N2)
	A/mallard/Ohio/81/1986(H6N2)
	A/mute swan/Rhode Island/A00325114/2008(H7N3)
	A/american green-winged
	A/northern
	shoveler/Mississippi/A00630207/2009(H7N6)
	A/chicken/Jiangxi/JXA132304/2013(H9N2) ⁻ A/mallard/South
	Dakota/A00536114/2007(H10N7)
	A/mallard/New York/A00919062/2009(H10N7)
	A/chicken/Jiangxi/JXA132717/2013 (H10N8) ^a
Avian	Newcastle disease virus (strain 2013 pt#3-3)
	Intectious bursal disease virus (strain P13- 03308)
	Infectious bronchitis virus (strain P14-00957ct)
	Reovirus (strain P14-00932)

Note: ^a: The RNA from these viruses was extracted directly from a clinical swab sample; for all others viruses, the RNA was extracted from virus isolates.