

Electronic Supplementary Material

Genomic Characterization of the First Parechovirus in Bats

Chang Zhang^{1,2#}, Jinliang Sheng^{3#}, Zihan Zhao², Chao Yan², Changchun Tu^{1,2,4}, Biao He^{2,4,✉}

1. College of Animal Science, Fujian A&F University, Fuzhou 350002, China
2. Key Laboratory of Jilin Province for Zoonosis Prevention and Control, Military Veterinary Institute, Academy of Military Medical Sciences, Changchun 130122, China;
3. College of Animal Science and Technology, Shihezi University, Shihezi 832003, China
4. Jiangsu Co-innovation Center for Prevention and Control of Important Animal Infectious Diseases and Zoonosis, Yangzhou 225009, China.

Supporting information to DOI: 10.1007/s12250-019-00108-1

Table S1 Primers for PCR analysis. Primers for BtPeV-F1, BtPeV-F2, and BtPeV-R were used to screen bat parechovirus in samples, the rest were used for whole-genome amplification. Locations of the primers refer to their position in the genome of BtPeV QAPp32 (MK348056)

Primer	Sequence (5'→3')	Polarity	Location
BtPeV-F1	GCAATGGCTCGCCTGTTT	+	2126-2143
BtPeV-F2	AGGACTTGCCGATTTTGG	+	2230-2247
BtPeV-R	CAGTTGCCAAGACATCAGG	-	2607-2625
5' RACE-R1	GATTACGCCAAGCTTGCCCAATTTTCCTTCCTGGACTGG	-	584-606
5' RACE-R2	GATTACGCCAAGCTTGAATGTGTAGTCTCTATACACAC	-	657-679
F1	CARAAYTCAGATMGAATGGC	+	416-435
R1	CGAACGGCACGAGAAACGAA	-	2583-2602
F2	GCGGATGCTGCTGCTGAGAACA	+	1883-1904
R2	GCCCCATCGGGGTAGAGTAAAT	-	5313-5334
F3	GGAAGACATCTGTGCAAAGC	+	4634-4653
R3	AGGTCCAGTGCACCAACAAT	-	6741-6760
F4	CTGACAATGTGCCTAAGGTT	+	6661-6680
R4	TGCARRACYAGCTCATACTC	-	6827-6846
3'RACE-F	GATTACGCCAAGCTTGAAGTGCATCTGGGTCTGACTG	+	6616-6638