

Electronic Supplementary Material

Drug resistance mutation profiles of the drug-naïve and first-line regimen-treated HIV-1-infected population of Suzhou, China

Tingting Li^{1,2#}, Feng Qian^{3#}, Ting Yuan¹, Weilu Xu³, Li Zhu³, Jinlong Huang³, Haiyan Wang³, Yueping Zhu³, Yinling Wang³, Xiaohong Li³, Saihong Gu³, Zhuqing Tan³, Hui Chen³, Xiangrong Luo³, Wei Zhu³, Wenjuan Lu³, Ping Xu³, Ming Li³, Yuying Chen^{1,2}, Yong Gao⁴, Rongge Yang¹, Chuanwu Zhu^{3✉}, Binlian Sun^{1✉}

1. Research Group of HIV-1 Molecular Epidemiology and Virology, The State Key Laboratory of Virology, Wuhan Institute of Virology, Chinese Academy of Sciences, Wuhan 430071, China
2. University of Chinese Academy of Sciences, Beijing 100000, China
3. Division of HIV-1/AIDS, the Fifth People's Hospital of Suzhou, Suzhou 215000, China
4. Department of Microbiology and Immunology, Schulich School of Medicine & Dentistry, University of Western Ontario, London N6A 5C1, Canada

Supporting information to DOI: 10.1007/s12250-017-4002-y




Figure S1. Recombination breakpoint analysis of the 13 URFs. Bootscan plot of percentage permuted trees over nucleotide position across the lengths of RT sequences.