## **Electronic Supplementary Material**

## Potential m6A and m5C Methylations within the Genome of A Chinese African Swine Fever Virus Strain

Lijia Jia<sup>1,2,7</sup>, Jianjun Chen<sup>1,2,3,6</sup>, Haizhou Liu<sup>2</sup>, Wenhui Fan<sup>4</sup>, Depeng Wang<sup>5</sup>, Jing Li<sup>4,7⊠</sup>, and Di Liu<sup>1,2,3,6,7⊠</sup>

 CAS Key Laboratory of Special Pathogens and Biosafety, Wuhan Institute of Virology, Chinese Academy of Sciences, Wuhan 430071, China
Computational Virology Group, Center for Bacteria and Viruses Resources and Bioinformation, Wuhan Institute of Virology, Chinese Academy of Sciences, Wuhan 430071, China
African Swine Fever Regional Laboratory of China, Wuhan 430071, China
CAS Key Laboratory of Pathogenic Microbiology and Immunology, Institute of Microbiology, Chinese Academy of Sciences, Beijing 100101, China
NextOmics Biosciences, Wuhan 430074, China
Center for Biosafety Mega-Science, Chinese Academy of Sciences, Wuhan 430071, China
University of Chinese Academy of Sciences, Beijing 100049, China
Supporting information to DOI: 10.1007/s12250-020-00217-2

Table S1 Methylation sites predicted with nanopolish and deepmod that matched the prediction of Tombo.

www.virosin.org