

Complete Genomic Sequence of Transmissible Gastroenteritis Virus TS and 3' End Sequence Characterization Following Cell Culture*

Jian-qiang LI^{1,2}, Jie CHENG¹, Xi LAN¹, Xue-rui LI¹, Wei LI², Xiang-ping YIN¹, Bao-yu LI¹,
Bin YANG¹, Zhi-yong LI¹, Yun ZHANG¹ and Ji-xing LIU^{1**}

(1. Key laboratory of Animal Virology of Ministry of Agriculture, State Key Laboratory of Veterinary Etiological Biology, Lanzhou Veterinary Research Institute, Chinese Academy of Agricultural Sciences, Lanzhou 730046, China; 2. Lanzou Institute of Biological Products, Lanzhou 730070, China)

Abstract: The complete genome sequence of transmissible Gastroenteritis virus (TGEV) strain TS, previously isolated from Gansu province, was cloned and compared with published sequence data from other TGEV strains. Phylogenetic tree analysis based on the amino acid and nucleotide sequences of the *S* gene showed that the TGEV strains were divided into 3 clusters. TGEV TS showed a close evolutionary relationship to the American Miller cluster but had a 5' non-translated region (NTR) sequence closely related to the American Purdue cluster. Continued culture in different cell types indicated that TGEV TS virulence could be attenuated after fifty passages in Porcine kidney (PK-15) cells, and that the Porcine kidney cell line IB-RS-2 (IBRS) was not suitable for culture of the TGEV strain TS.

Key words: Cloning; Complete genome; Gastroenteritis virus (TGEV); Cell culture

Transmissible gastroenteritis virus (TGEV) causes acute infection in swine of all ages, and the disease is characterized by severe enteritis and diarrhea. These symptoms are especially severe in newborn piglets less than 2 weeks old, frequently leading to death with mortality rates up to 100% [9, 15, 17]. TGEV was first reported in the United States in 1946 and has now been reported in many other countries, including

China [10, 23, 24]. TGEV is now prevalent throughout China and is a cause of serious economic losses.

TGEV is a member of the *Coronavirus* genus and the *Coronaviridae* family. Its genome consists of a single molecule of positive-sense, single stranded RNA, 28.5 kb in size, which is transcribed into several subgenomic mRNAs for the production of four structural proteins [Spike protein (S), membrane protein (M), small membrane protein (E) and nucleocapsid protein (N)] and four non-structural proteins (Replicase, 3a, 3b, and Protein 7) [16]. The genes are arranged in the order 5'-*rep-s-3a-3b-e-m-n-7-3'* [21]. Transcription regulatory sequences (TRSs) including a

Received: 2009-10-29, Accepted: 2010-04-14

* Foundation items: National Basic Research Program (2004CCA00500); National High-tech Development Research Program of China (2006AA02Z440).

** Corresponding author.

Phone: +86-931-8342682, E-mail: liujixing@hotmail.com

highly conserved core sequence (CS), 5'-CUAAAC-3', or a related sequence, are present at sites immediately upstream of the TGEV genes^[1]. The S glycoprotein makes up the large surface projections of the virion and plays an important role in the attachment of viral particles to host cell receptors, with subsequent penetration into the cell by membrane fusion. The S glycoprotein also stimulates the production of neutralizing antibodies in the host^[18]. The M and E proteins are essential for viral envelope formation and release; the M protein can also stimulate the production of interferon (IFN)^[12]. The N protein participates in transcription of the viral genome, the formation of the viral core, and packaging of viral RNA^[20]. The Coronavirus replicase is a multifunctional polyprotein with helicase and protease activities necessary for the transcription of viral RNA^[4, 23]. Four major antigenic sites, A, B, C and D, were characterized on the N terminus of the S protein. Using monoclonal antibodies, the adjacent sites A and B were mapped to a region of approximately 200 amino acids (aa) beginning from residue 506 at its N terminal boundary. The two antigenic sites also overlapped with the S protein domain encompassing aa 522–744 that mediates aminopeptidase N (APN) receptor binding^[5, 6, 8]. After continuous passage in cell culture, TGEV isolates gradually lose their virulence and may shift from enteric to respiratory tropism^[14]. Attenuation of virulence and tropism shift can also occur in nature, an example being the naturally occurring *s* and *orf3* gene deletion mutant, the Porcine respiratory coronavirus (PRCV), which has both reduced pathogenicity and a predominantly respiratory tropism^[14]. Although it is generally accepted that

deletions in the spike and *orf3* genes contribute to tropism change and attenuation in PRCV, other genes may also be involved. For instance, amino acid mutations in the M protein affect the ability of attenuated Purdue TGEV P115 to induce IFN- α , implying a potential role for the M protein in altered host response and virulence^[12].

In this study, the complete genome of TGEV TS strain isolated in Gansu province has been cloned. In addition, the virus was continuously propagated in different cell lines and the 3' end genome sequences were also cloned. These data are useful for further study of the molecular structure of the TGEV strains prevalent in China, and the characterization of the long-term TGEV genome stability in different host cells.

MATERIALS AND METHODS

Viruses and Cells

TGEV strain TS was isolated from a suburb in Gansu province, China. Swine testicle (ST), PK-15 and IBRS cells were grown as monolayers in DMEM (GIBCO, USA) containing 10% fetal calf serum (GIBCO, USA) and 5% CO₂ in air. Samples were obtained by passaging the first TGEV field isolate 10 times in swine intestine and named TS-ST.

RNA Extraction

Total RNA was isolated from infected cell samples using a RNA extraction kit (Qiagen, Germany) following the manufacturer's instructions.

RT-PCR

The RT-PCR amplifications were carried out using seventeen primer sets (Table 1) and an RT-PCR amplification Kit (Toyobo, Japan) according to the manufacturer's instructions (Fig. 1).

Table 1. Primers used in cloning the full-length TGEV TS genome

Primers	Position ^a	Sequence (5'—3')	Target gene	length ^b
A1S	1	ACTTTTAAAGTAAAGTGAGTGTAGC	5'NTR and Replicase <i>1a</i>	2957
A1R	2957	GTCACCACCACCCATTTTAT		
A2S	2933	GTATAATAAAATGGGTGGTG	Replicase <i>1b</i>	2053
A2R	4985	AGCATCACGGAAACCATAAT		
A3S	4948	ATATAGATTGGCAAGCTCAT		
A3R	7476	CACTTTTTGGAGACACAGAT		
A4S	7405	CATTTAATGCTGCTGTAGGTTCA		
A4R	10069	ACACTAGGAAGGACAAAAGA		
A5S	10044	TTTATGTCTTTTIGCCTTCC		
A5R	12385	TACTAACATGGTCTGGATCA		
B1S	12352	TAGTGCAGCTCGACTAGAAC		
B1R	14622	TAACCTAAATCCGCATAATC		
B2S	14610	GTGCTACAACAAAGATTATG	2599	
B2R	17208	GGTACATTAGTTCCAACATT	2230	
B3S	17176	TGTCTGTGGGATAATGTTG		
B3R	19405	TGCCTTACAATCTACAATGA		
B4S	19321	CTTGGACGATTTTGTGACTA		
B4R	20395	TACGACCAAACCACAAATA	1075	
S1S	20295	AAGGAAGGGTA GTTGCTCA	<i>s</i> gene	1267
S1R	21561	TACCGTGGTCCATC GTTAC		
S2S	21430	GTGCCACAGTGTTCATTG	2372	
S2R	23801	GCATCAGCACTCAATTCGTCAAGCCT		
S3S	23631	ACATCACGA GGTCTTGCTACT		
S3R	24872	GCACAATCA AGTTCGTCA AG	1432	
CS	24616	GCTGTGGATGCATAGGTTGT		
CR	26047	GTGCTCCATCGGGTTGTAT	<i>sM</i> gene	368
SMS	25771	GCGGTTCTAAACGAAATTGAC		
SMR	26138	CATGCAATCACACACGCTAAT	<i>m</i> gene	929
MS	25955	GTTGCAATTTAGGAAGGACAG		
MR	26883	TTGGCCATTTAGAAGTTTAGT	<i>n</i> gene	1215
NS	26821	GAACTGATAATTTGAGTGAGC		
NR	28035	GCATCTCGTTTAGTTCGTTAC	<i>orf7</i> and 3'NTR	561
TS	27988	ATGACACACAGGTTGAGATA		
TR	28548	TTTTTTGTATCACTATCAAA		

^a Position is relative to the TGEV TS strain; ^b Length of PCR products

Cloning and sequencing

RT-PCR products were purified using a Gel Extraction Mini Kit (TaKaRa) and were ligated to the pMD18-T plasmid (TaKaRa). Competent *E. coli* JM109 were transformed with the recombinant vector.

Positive clones were sequenced by the TaKaRa Biotechnology (Dalian) Co. Ltd.

Phylogenetic analysis

Phylogenetic analysis was performed for the TGEV TS strain compared to other coronavirus strains retrieved




Fig. 1. Strategies for sequence determination of the TGEV TS

from GenBank. The sequencing data were aligned using the Clustal W method (Lasergene program version 7.1, DNASTAR Inc. Madison, WI). Phylogenetic trees were constructed for the *s* and replicase genes using the bootstrap N-J method in the DNASTAR program using 1000 bootstraps. The following sequences, taken from GenBank, were included in the analysis: Feline infectious peritonitis virus (FIPV, AY994055), Porcine epidemic diarrhea virus (PEDV, AAK38661), Human coronavirus 229E (HCoV 229E, Q05002), Severe acute respiratory syndrome virus (SARSV, AAP41036), Avian infectious bronchitis virus (AIBV, P27920), Murine coronavirus (MCoV, P16342), Bovine coronavirus (BCoV, Q91A29), Porcine respiratory coronavirus (PRCV ISU-1, DQ811787), TGEV/Miller M6 (DQ811786), TGEV/Miller M60 (DQ811786), TGEV/Virulent Purdue (DQ811789), TGEV/Pur46-MAD (NC002326), TGEV/SC-Y (DQ443743), TGEV/Purdue P115 (DQ81178/US), TGEV/attenuated H/ EU074218/ China, TGEV/Purdue P-15 (X06371), TGEV/HN-2002 (AY587882), TGEV/FS772-70 (X53128), TGEV/TSX (DQ001167), TGEV/96-1933 (AF104420), TGEV/TO14 (AF302263), TGEV/TH-98 (AF494337).

Characterization of TS after continuous passage in cultured cells

TS was plaque-purified and passaged in ST, PK-15 and IBRS cells 50 times. The clones were named

TS-ST₅₀, TS-PK₅₀ and TS-IB₅₀ respectively. The 3' end sequences of the TS-ST₅₀, TS-PK₅₀ and TS-IB₅₀ genomes were amplified and cloned.

RESULTS

Complete genome sequence of the TGEV TS strain

The complete genome of the TGEV TS strain comprised 28 541 nucleotides (nt), excluding the poly (A) tail. The 5' end two-thirds of the genome nt 1-20368 contained a 314 nt non-translated region (NTR) and an open reading frame (ORF) encoding the viral RNA-dependent RNA replicase. The remaining 8 173 nt incorporated the structural genes, *orf3* and *orf7*, and a 3'NTR. A small ORF (*uorf*) with the potential to code for a 3-amino acid peptide at the 5' NTR of the genome, from nucleotide position 99-108 (5'-ACUU UAAAAG-3'), was also found. A conserved sequence was present at different distances from the initiation codon of each gene, and the 3' end of the TGEV genome had a poly (A) tail of undetermined length. The complete TGEV TS genome sequence has been submitted to NCBI and has been given the GenBank Accession Number, DQ201447.

The non-structural genes

The replicase genes were designated genes *la* and *lb*, and share 43 overlapping nucleotides. The replicase *la* gene was predicted to encode a protein of 4017 aa

Table 2. Length in amino acids of predicted structural and nonstructural proteins of the seven TGEV isolates and PRCV-ISU-1

	TGEV								PRCV
	TS	Miller M6	Mille M60	H	SC-Y	Pur46-MAD	Virulent Purdue	Purdue P115	ISU-1
Replicase 1a	4017	4017	4017	4017	4017	4017	4017	4017	4014
Replicase 1b	2680	2680	2680	2680	2680	2680	2680	2680	2680
S	1449	1449	1448	1448	1447	1447	1449	1447	1222
3a	72	72	72	72	71	71	71	71	0
3b	244	244	67	244	244	244	244	244	205
E	82	82	82	82	82	82	82	82	82
M	262	262	264	262	262	262	262	262	261
N	382	382	382	382	382	382	382	382	382
7	78	78	78	78	78	78	78	78	78

while the replicase *1b* gene was predicted to encode a protein of 2 680 aa. Nucleotide sequence analysis revealed that there were no deletions or insertions in the replicase genes of all TGEV strains. Compared with the TGEV sequences, PRCV ISU-1 encoded a replicase 1b protein of identical length and a replicase 1a with a 3 amino acid deletion (Table 2).

TGEV TS strain *orf3a* and *orf3b* genes were predicted to encode proteins of 72 aa and 244 aa, respectively, identical to the corresponding Miller M60 genes. Compared with other TGEV strains, a 3 nt insertion present in the TS, as Miller M6 and Miller M60 strains led to one amino acid (histidine) being inserted at the end of *orf3a*. No deletions or insertions were found in the TS *orf3b* gene compared to other strains except for Miller M60. A 531 nt deletion in the *orf3b* gene of Miller M60 resulted in a truncated 3b protein of 67 aa in length. Meanwhile, a 184 nt deletion in the *orf3a* gene of PRCV-ISU-1 disrupted the predicted open-reading frame of its encoded protein, and a 117 nt deletion in the *orf3b* gene predicted a shorter nonstructural protein 3b compared to other TGEV strains. The TGEV TS strain *orf7* gene was predicted to encode a protein of 78 amino acids, and no insertions or deletions were found compared

with other TGEV strains and PRCV ISU-1 (Table 2).

The structural genes

The nucleotide sequence of the TGEV TS strain *s* gene was 4 350 nt in length, encoding a predicted protein of 1 449 amino acids. As such, it was identical in length to the equivalent proteins of the Miller M6 and virulent Purdue strains. Compared with the TGEV M6 strain, a 3 nt deletion was found in the *s* gene of Miller M60 at position 2 385 - 2 387, which predicted a spike protein of 1 448 amino acids in length. A 6 nt deletion in the *s* genes of Pur46-MAD and Purdue P115 predicted that their spike proteins were 2 amino acids shorter than that of the TGEV TS strain. Sequence analysis confirmed the previous report of a 681 nt deletion in the 5' end of the *s* gene of PRCV-ISU-1^[14].

There were no deletions or insertions in the *e* and *n* genes of all the TGEV strains. The deduced amino acid sequences of the *e* and *n* genes of TGEV and PRCV ISU-1 were 82 and 382 amino acids in length, respectively. The *m* gene of M60 had a 6 nt insertion compared with other TGEV strains, predicting a membrane protein 2 amino acids longer than in the other strains. PRCV-ISU-1 had a 3 nt deletion in the *m* gene compared with TGEV TS strain, predicting an M

Table 3. The nucleotides and amino acids homology of genomes compare the TGEV TS to other TGEV strains and PRCV-ISU-1 (%)

Virus	ORF1		S		ORF3a		ORF3b		sM		M		N		ORF7	
	nt	aa	nt	aa	nt	aa	nt	aa	nt	aa	nt	aa	nt	aa	nt	aa
Miller M6	99.8	99.7	99.9	99.6	99.1	97.3	99.6	98.8	99.6	98.8	99.7	99.2	99.9	99.7	99.6	100
Miller M60	99.8	99.6	99.8	99.5	98.6	95.9	98.5	95.7	99.6	98.8	99.7	99.2	99.7	99.5	99.6	98.8
H attenuated	99.7	99.6	99.6	99.0	98.6	95.6	99.5	98.4	98.8	96.4	99.6	98.9	99.8	99.7	100	100
SC-Y	98.6	97.6	98.2	97.8	92.7	87.8	98.6	96.7	98.8	96.4	98.0	97.0	98.0	97.9	96.2	93.8
Pur46-MAD	98.9	98.1	98.5	98.2	96.8	87.8	98.5	96.3	98.8	96.4	98.1	97.3	98.1	98.2	96.2	93.8
Virulent Purdue	99.0	98.2	98.6	98.6	96.8	87.8	98.8	97.2	99.2	97.6	98.2	97.7	98.2	97.9	96.2	93.8
Purdue P115	98.8	98.0	98.4	98.1	96.8	87.8	98.6	96.7	98.8	96.4	98.1	97.3	98.2	98.4	96.2	93.8
PRCV ISU-1	98.0	96.8	97.7	98.4	---	---	97.1	93.7	98.4	100	98.1	97.3	97.4	98.2	97.9	100

protein of 261 amino acid in length.

Homology comparisons

The nucleotide and predicted amino acid sequences of each region (replicase, S, 3a, 3b, E, M, N and ORF7) of TGEV TS, other TGEV reference strains and PRCV ISU-1 were compared. Protein N exhibited the highest amino acid similarity (97.9%-99.7%), protein E showed the lowest amino acid similarity (96.4% - 100%), while the amino acid homologies of proteins S and M were 97.8% - 99.6% and 97.3% - 99.2%, respectively. In the non-structural protein regions, the replicase was highly conserved (97.6% - 99.7% amino acid similarity), while the 3b (95.7% - 98.8% amino acid similarity) and ORF7 (93.8% - 100% amino acid similarity) proteins were less

conserved. The highly variable protein 3a showed the lowest conservation (87.8% - 97.3% amino acid similarity). TGEV TS ORF7 shared 100% homology, both in nucleotide and amino acid sequences, with the attenuated strain H also isolated in China (Table 3).

Phylogenetic tree analysis

In order to determine the relationship between the TGEV TS strain and other Coronaviruses, a phylogenetic tree based on the Coronavirus replicase amino acid sequences was constructed. The data indicated that all TGEV strains and PRCV ISU-I clustered into one group, and that the TGEV strains were most closely related to FIPV. SARS, which was discovered in China in 2003, had a closer relationship with MHV and BCoV, which belong to the Coronavirus




Fig. 2. Phylogenetic tree with bootstrap N-J method constructed based on the TGEV strains and other Coronavirus replicase amino acid sequences.

group 2 (Fig. 2). The tree based on the deduced amino acid sequences of the TGEV replicase genes revealed that the strains compared were divided into 2 groups: the Miller group and the Purdue group (Fig. 3). The tree based on the nucleotide and amino acid sequences of the TGEV *s* genes showed that all the TGEV strains analyzed were divided into 3 clusters, and that the TGEV TS strain shared the closest relationship with Miller M6 and HN2002 (Fig. 4). Another phylogenetic tree based on the nucleotide sequences of the *s* genes of TGEV strains and PRCV ISU-1 was also constructed. This revealed that all TGEV strains analysed, except 96-1933, were divided into 3 clusters (Fig. 5). All phylogenetic trees demonstrated that the

TGEV TS strain shared the closest relationship with Miller M6, while PRCV was most closely related to TSX of the British group of coronaviruses.

Characterization of TS genome stability in long-term cell culture

After passaging in swine testicle (ST) cells for 10 generations, cytopathic effects (CPE) were observed after 48 h in culture, and the genes could be cloned from the cells that displayed CPE. CPE was also observed 48 - 72 h from the start of continued passage in PK-15 and IBRS cells for 10 generations, but the genes could not be cloned from the cells that displayed CPE. After passage for 50 generations, the 3' ends of the viral genes could be cloned from ST and




Fig. 3. Phylogenetic tree with bootstrap N-J method was constructed based on the TGEV TS and other TGEV strains replicase amino acid sequences.




Fig. 4. Phylogenetic tree with bootstrap N-J method was constructed based on the TGEV TS and other TGEV strains S amino acid sequences.




Fig. 5. Phylogenetic tree with bootstrap N-J method was constructed based on the TGEV TS and other TGEV strains *s* gene nucleotide sequences.

Table 4. The *s* and *orf3* nucleotides mutations from TS to TS-ST₅₀ and TS-PK₅₀

	Position	TS	TS-ST	TS-PK
<i>s</i> gene	158	T	T	C
	202	G	T	T
	290	G	C	C
	295	C	C	T
	952	T	T	C
	1253	T	T	T
	1691	C	C	A
	1753	T	T	G
	3629	T	C	C
<i>orf3</i>	50	G	A	A
	85	G	G	A
	176	C	T	T
	660	T	T	G
	910	A	A	G
	998	G	G	G

PK-15 CPE cells, but the genes could not be amplified from IBRS cells that displayed CPE.

The RT-PCR results from ST (TS-ST₅₀) and PK-15 (TS-PK₅₀) cells cultured with TS for 50 passages indicated that there were no nucleotide deletions or insertions at the 3' end of the TGEV genome. However, there were nucleotide mutations in the *s* and *orf3* genes, with TS-ST₅₀ showing less variation than TS-PK₅₀ (Table 4).

DISCUSSION

Eight complete nucleotide sequences of the TGEV

genome have been reported in GenBank, of which three were collected in China. Analysis of the full genomic sequence of the TGEV TS strain can contribute to the understanding of the genetic structure, diversity, evolution and, in particular, the prevalence of these coronaviruses in China.

The uORF is present in all Coronaviruses and plays an important role in virus replication and packaging. The uORF sequences of HCoV-299E and IBV were found to be eleven codons in length, while those of MHV and PEDV were eight and twelve codons, respectively. Meanwhile, the uORF of TGEV encodes a peptide of only three-amino acids^[11]. In this study, we demonstrated that the TS strain has the same three-amino acid uORF peptide as other TGEV strains, indicating that uORF is highly conserved in TGEV. Moreover, most of the 5' NTR sequences containing uORF were conserved, except for the first few nucleotides. In this study, we used two sense PCR primers beginning with the sequences "ACTTTTAAA GTAAAGTGAGT" and "GCCAGTAAAGTGAGTG TAGC", which corresponded to the Pur46-MAD and Miller M6 sequences, respectively. The PCR results showed that only the former primer could amplify the

5' NTR of the TGEV TS strain, indicating that the TGEV TS strain M6 sequence should start with the "ACTTTTAAAGTAAAGTGAGT" sequence found in the Pur46-MAD strain. This was confirmed by 5' RACE (data not shown). Alignment of all TGEV sequences showed that they were highly conserved except for nucleotide deletions in some strains (Fig. 6). We speculate that incomplete sequencing is responsible for the presence of these deletions, and propose that the first several nucleotides of the 5' NTR are conserved among the TGEV strains. The length of the poly (A) tail of the TGEV genome was not confirmed because it has not been possible to amplify it accurately by PCR, although a poly(A) sequence of 9 residues has been cloned by 3' RACE (data not shown). Whether this sequence corresponds to the true poly (A) tail remains unknown. Nevertheless, minigenomes and full length infectious RNAs with a poly(A) tail of 24 residues have been constructed and replicate efficiently, indicating that 24 A residues are sufficient for TGEV RNA replication [2].

The principal component of the Coronavirus genome is the replicase gene, which contains two large open reading frames, *orf1a* and *orf1b*, from which two products are translated. The latter is synthesized via a ribosomal frameshifting mechanism, facilitated by a pseudoknot structure [7]. *orf1b*

sequences in two Chinese TGEV strains, TS and SC-Y, were predicted to encode a protein of 2 678 aa. In fact, because the ribosomal frameshifting mechanism was not accounted for, two amino acids were deleted in ORF1b of TS and SC-Y. In common with other TGEV strains, the predicted *orf1a* sequences of TS and SC-Y extended from nucleotide 315 to 12 054. This resulted in a 4 017-codon *orf* that overlapped *orf1b* from nucleotide 12 011 to nucleotide 20 368, with the capacity to encode a polypeptide of 2 680 amino acids (Fig. 7).

The *orf3* and *s* genes have been hypothesized to be important in the virulence and pathogenesis of TGEV infections [19]. Analysis of TGEV ORF3 sequences revealed that TS had equal numbers of ORF3 nucleotides and amino acids as the virulent Miller M6 strain. The G residue at nucleotide position 655 of the spike protein was essential to maintain enteric tropism of the TGEV strain PUR46-MAD, and mutation at this nucleotide caused a shift from enteric to respiratory tropism [3]. The T to G mutation at nucleotide 1 753 of the *s* gene is regarded to be a signal of attenuation [25]. Analysis of the *s* gene sequence of the TGEV TS strain revealed a G at nucleotide 655 and a T at nucleotide 1 753 indicating that TS was a virulent strain, also demonstrated by animal experiments (data not shown). A 6 nt (nt 1 122




Fig. 6. 5'NTR sequence alignment of TGEV strains and PRCV ISU-1.

```

11994 AGTTC TACTGTTGATCAAAGTTATTTAAACGAGTGC GGGGTTTTAGTGCAGCTCGACTAG      12054 ORF1a.
      S  S  T  V  D  Q  S  Y  L  N  E  C  G  V  L  V  Q  L  D  .  ↵
11994 AGTTC TACTGTTGATCAAAGTTATTTATTTAAACGAGTGC GGGGTTTTAGTGCAGCTCGACTAGAA 12060 ORF1ab.
      S  S  T  V  D  Q  S  Y  L  F  K  R  V  R  G  F  S  A  A  R  L  E  ↵
11994 AGTTC TACTGTTGATCAAAGTTATTTAaagttatttaAAACGAGTGC GGGGTTTTAGTGCAGCTCGACTAGAA 12066 ORF1ab.
      S  S  T  V  D  Q  S  Y  L  K  L  F  K  R  V  R  G  F  S  A  A  R  L  E  ↵

```

Fig. 7. Nucleotide and amino acid sequence of the ribosomal frameshifting region. The first two line is the nucleotides and a.a. of ORF1a, the second two line is the nucleotides and predicted a.a. of ORF1ab that had been in the GenBank, the third two line is the nucleotide and predicted aa of ORF1ab of ribosomal frameshifting, the nucleotides with underline is the ribosomal binding site, the nucleotides with box is the insertion of ribosomal frameshifting.

to 1 127, aa 375 - 376) deletion is present in the attenuated TGEV Purdue P115 and Pur46-MAD spike genes compared to the virulent Purdue strain, and so this deletion was also thought to play a role in attenuation^[18]. Comparison of the attenuated Miller M60 strain with the virulent Miller M6 strain showed that no deletion was present at nucleotides 1 122 to 1 127, but there was a deletion at nucleotides 2 386-2 388 in the *s* gene. This suggested that the deletion in the *s* gene may affect virus virulence, but this is uncertain from the position of this particular deletion. The TGEV TS S protein possessed the same number of amino acids as the Miller M6 and virulent Purdue strains, suggesting that TS is also a virulent strain. Amino acid mutations in the M protein affect its ability to induce IFN- α , and so the M protein has a potential role in TGEV virulence. The *m* gene has been regarded as a highly conserved TGEV gene, but comparison of the lengths of TGEV *m* genes revealed that the Miller M60 gene (789 nucleotides) encoded a 263 aa polypeptide, the TGEV gene encoded a 262 aa polypeptide, and the PRCV ISU-1 gene encoded a 261 aa polypeptide. The aa insertion in TGEV is suggestive of the M protein being subject to environmental selection pressure because it affects a

coat protein. Although the ORF7 nucleotide deletion of TGEV had been considered to affect virulence^[13], we found no deletion among the TGEV strains and PRCV ISU-1.

Homology and phylogenetic tree analysis showed that TGEV TS had the closest sequence relationship to the Miller M6 strain. All the compared TGEV strains were divided into 3 clusters in the phylogenetic tree based on the *s* gene nucleotide and amino acid sequences. These were the American Miller cluster, American Purdue cluster and British cluster. The attenuated strain H from China had the closest relationship with Miller M60, while another Chinese strain, SC-Y, belonging to the Purdue cluster was separated into an independent group based on the S amino acid phylogenetic tree. Analysis of SC-Y gene revealed that one nucleotide mutation generating a termination codon appeared at aa residue 924. This may have been due to a sequencing mistake because no such mutations have been described in coronavirus *s* genes. Surprisingly, the *orf3a* genes of viruses in the Miller cluster were 72 aa in length, while in the Purdue cluster they were 71 aa in length. Therefore, the nucleotide number of the *orf3a* genes may differentiate the Miller cluster and Purdue cluster

strains. Homology analysis indicated that PRCV ISU-I had the closest relationship with the Miller strains at both the nucleotide and protein levels. The *orf3* gene of the Miller strains also had a high frequency of mutations and deletions, suggesting that a Miller-like strain was the ancestor of PRCV-ISU-1 [25]. In this study, we confirmed that PRCV-ISU-1 had a closer relationship with Miller strains than the Purdue strains. The TS strain shared 100% aa homology with PRCV-ISU-1 in the *e* gene, indicating that TS may have the closest relationship with the PRCV-ISU-1. Interestingly, the phylogenetic tree based on the *s* gene sequence showed that PRCV-ISU-1 had the closest relationship with TSX, which belongs to the British cluster.

As described above, we have determined the entire genomic sequence of the TGEV TS strain isolated from the Gansu province of China. We characterized the TS strain by following its passage in three different cell types typically used for the culture of TGEV. As expected, ST cells were most sensitive to the presence of the TGEV virus, as indicated by the presence of cytopathic effects (CPE) after 48 h in culture. Although CPE were obvious in IB cells passaged for 50 generations, TGEV genes could not be cloned by RT-PCR, suggesting that maybe the virus can not be propagated in the cells. This indicated that IB cells were not suitable for the culture of the TGEV virus. The 3' end genome sequences of TS-ST₅₀ and TS-PK₅₀ showed no nucleotide mutations in the *m*, *n*, *e* and *orf7* genes. However, the *s* and *orf3* genes of TS-PK₅₀ had more nucleotides mutations than in TS-ST₅₀. Surprisingly, the T residue at position 1753 of the TS *s* gene mutated to G in TS-PK₅₀. This is regarded as a signal of attenuation, indicating that

TGEV TS virulence could be attenuated by continued passage in PK-15 cells.

Acknowledgement

The financial support of this work was provided by grants from the National basic research program (2004CCA00500) and National high-tech development research program of China (2006AA02Z440).

References

1. **Alonso S, Izeta A, Sola I, et al.** 2002. Transcription regulatory sequences and mRNA expression levels in the coronavirus transmissible gastroenteritis virus. *J Virol*, 76: 1293-1308.
2. **Ander I, Cristian S, Sara A, et al.** 1999. Replication and packaging of transmissible gastroenteritis coronavirus-derived synthetic minigenomes. *J Virol*, 73: 1535-1545.
3. **Ballesteros M L, Sanchez C M, Enjuanes L.** 1997. Two Amino Acid Changes at the N-Terminus of Transmissible Gastroenteritis Coronavirus Spike Protein Result in the Loss of Enteric Tropism. *Virology*, 227: 378-388.
4. **Bredenbeek P J, Pachuk C J, Noten A F, et al.** 1990. The primary structure and expression of the second open reading frame of the polymerase gene of the coronavirus MHV-A59; a highly conserved polymerase is expressed by an efficient ribosomal frameshifting mechanism. *Nucl Acids Res*, 18: 1825-1830.
5. **Correa I, Jimenez G, Sune C, et al.** 1988. Antigenic structure of the E2 glycoprotein from transmissible gastroenteritis coronavirus. *Virus Res*, 10: 77-93.
6. **Delmas B, Rasschaert D, Godet M, et al.** 1990. Four major antigenic sites of the coronavirus transmissible gastroenteritis virus are located on the amino-terminal half of spike glycoprotein S. *J Gen Virol*, 71: 1313-1323.
7. **Eleouet J F, Rasschaert D, Lambert P, et al.** 1995. Complete Sequence (20 Kilobases) of the Polyprotein-Encoding Gene 1 of Transmissible Gastroenteritis Virus. *Virology*, 206: 817- 822.
8. **Godet M, Grosclaude J, Delmas B, et al.** 1994. Major receptor-binding and neutralization determinants are located within the same domain of the transmissible gastroenteritis virus (coronavirus) spike protein. *J Virol*,

- 68: 8008-8016.
9. **Isabel C, Fatima G, Maria J B, et al.** 1990. Localization of antigenic sites of the E2 glycoprotein of transmissible gastroenteritis coronavirus. *J Gen Virol*, 71: 271-279.
 10. **Kemeny L J, Woods R D.** 1977. Quantitative transmissible gastroenteritis virus shedding patterns in lactating sows. *Am J Vet Res*, 38: 307-310.
 11. **Kocherhans R, Bridgen A, Ackermann M, et al.** 2001. Completion of the Porcine Epidemic Diarrhoea Coronavirus (PEDV) Genome Sequence. *Virus Genes*, 23: 137-144.
 12. **Woods R D, Wesley R D.** 1986. Immune response in sows given transmissible gastroenteritis virus or canine coronavirus. *Am J Vet Res*, 47: 1239-1242.
 13. **Ortego J, Sola I, Almazan F, et al.** 2003. Transmissible gastroenteritis coronavirus gene 7 is not essential but influences in vivo virus replication and virulence. *Virology*, 308: 13-22.
 14. **Saif L J, Sestak K.** 2006. Transmissible gastroenteritis virus and porcine respiratory coronavirus, In: *Diseases of Swine* (Zimmerman, J J, et al. Ed), 9th ed. Iowa: Iowa State University Press, p489-516.
 15. **Sanchez C M, Izeta A, Sanches-morgado J M, et al.** 1999. Targeted recombination demonstrates that the spike gene of transmissible gastroenteritis coronavirus is a determinant of its enteric tropism and virulence. *J Virol*, 73: 7607-7618.
 16. **Serge B, Hubert L.** 1995. Site-specific alteration of transmissible gastroenteritis virus spike protein results in markedly reduced pathogenicity. *J Gen Virol*, 76: 2235-2241.
 17. **Siddell S, Wege H, Meulen Ter V.** 1983. The Biology of Coronaviruses. *J Gen Virol*, 64: 761-776 .
 18. **Simkins R A, Sail L J, Weinau P A.** 1989. Epitope mapping and the detection of transmissible gastroenteritis viral proteins in cell culture using biotinylated monoclonal antibodies in a fixed-cell ELISA. *Arch Virol*, 107: 179-190.
 19. **Vaughn E M, Halbur P G, Paul P S.** 1995. Sequence comparison of porcine respiratory coronavirus isolates reveals heterogeneity in the S, 3, and 3-1 genes. *J Virol*, 69: 3176-3184.
 20. **Wang Y, Zhang X.** 1999. The Nucleocapsid Protein of Coronavirus Mouse Hepatitis Virus Interacts with the Cellular Heterogeneous Nuclear Ribonucleoprotein A1 in Vitro and in Vivo. *Virology*, 265: 96.
 21. **Woods R D.** 1976. Leukocyte-aggregation assay for transmissible gastroenteritis of swine. *Am J Vet Res*, 37: 1405-1408.
 22. **Woods R D, Wesley R D.** 1986. Immune response in sows given transmissible gastroenteritis virus or canine coronavirus. *Am J Vet Res*, 47: 1239-1242.
 23. **Yang H T, Xie W Q, Xue X Y, et al.** 2005. Design of wide-spectrum inhibitors targeting coronavirus main proteases. *Plos Biology*, 10: 1742-1752.
 24. **Yin J C, Ren X F, Li Y.** 2005. Molecular Cloning and Phylogenetic Analysis of ORF7 Region of Chinese Isolate TH-98 from Transmissible Gastroenteritis Virus. *Virus Genes*, 30: 395-401.
 25. **Zhang X S, Hasoksuz M, Spiro D, et al.** 2007. Complete genomic sequences, a key residue in the spike protein and deletions in nonstructural protein 3b of US strains of the virulent and attenuated coronaviruses, transmissible gastroenteritis virus and porcine respiratory coronavirus. *Virology*, 358: 424-435.