



Research Article

Insights into cross-species infection: Porcine epidemic diarrhea virus infections in the rodent



Jianing Chen^{a,1}, Zemei Wang^{a,b,1}, Shengyu Lin^{a,c}, Mengling Gao^a, Yongheng Shao^{a,b}, Shuxian Li^{a,d}, Qingbo Chen^{a,d}, Yaru Cui^a, Yonghao Hu^b, Guangliang Liu^{a,*}

^a State Key Laboratory for Animal Disease Control and Prevention, College of Veterinary Medicine, Lanzhou University, Lanzhou Veterinary Research Institute, Chinese Academy of Agricultural Sciences, Lanzhou 730000, China

^b College of Veterinary Medicine, Gansu Agricultural University, Lanzhou 730070, China

^c College of Animal Sciences, Fujian Agriculture and Forestry University, Fuzhou 350002, China

^d College of Veterinary Medicine, Xinjiang Agricultural University, Urumqi 830052, China

ARTICLE INFO

Keywords:

Porcine epidemic diarrhea virus

Rodent

Cerebral tropism

Cross-species

Animal model

ABSTRACT

The cross-species infection of coronaviruses has resulted in several major epidemics since 2003. Porcine epidemic diarrhea virus (PEDV) is a devastating swine enteric coronavirus, which targets pigs as the only natural reservoir. In this study, the nucleic acid of PEDV was detected in rat fecal samples collected from pig farms. Further animal tests showed that PEDV can cause systemic infections in neonatal mice and rats via intracranial inoculation. The brain, lung, intestine and spleen were all targets for PEDV in rodents in contrast to the intestine being targeted in pigs. Morbidity and mortality vary via different infection routes. PEDV was also detectable in feces after infection, suggesting that the infected rodents were potential infectious sources. Moreover, the cerebral tropism of PEDV was verified in piglets via orally inoculation, which had not been identified before. In conclusion, our findings demonstrate that PEDV could cross the species barrier to infect mice and rats through different routes in experimental settings. Although it is highly devastating to piglets, PEDV changes the target organs and turns to be milder when meeting with new hosts. Based on these findings, more attention should be paid to the potential cross-species infection of PEDV.

INTRODUCTION

Porcine epidemic diarrhea (PED) is a devastating enteric disease in pigs, characterized by severe enteritis, vomiting, and watery diarrhea. The causative agent, porcine epidemic diarrhea virus (PEDV), is an enveloped, single-stranded, positive-sense RNA virus belonging to the *Coronaviridae* family. PEDV can infect pigs of all ages, but morbidity and mortality vary. It can cause 100 % mortality in piglets within two weeks but is much milder in pigs older than two weeks. Since its first discovery in the 1970s (Wood, 1977; Pensaert and de Bouck, 1978), PEDV has subsequently spread worldwide. In the 2010s, variant PEDV strains re-emerged, eliminating millions of suckling piglets (Sun et al., 2012; Stevenson et al., 2013; Chung et al.,

2016; Sun et al., 2019). To date, the virus is still evolving and circulating worldwide.

Pigs are identified as the only natural reservoir for PEDV. However, this virus exhibits poor replication in swine cells. Both IPEC-J2 and IPI-2I, two swine intestine cell lines, are permissive for PEDV infection but with limited viral replication (Zhao et al., 2014; Wang et al., 2019). In contrast, cell lines derived from other species are susceptible to PEDV. The rat crypt epithelial cell line (IEC-6), Vero cells, 293 cells, various human hepatocyte cell lines and human intestinal cell lines support PEDV replication to high titers (Zhang et al., 2017; Chen J. et al., 2020; Lv et al., 2022; Niu et al., 2023). Recently, several PEDV-like coronaviruses were also identified by virome analysis in bats in China, Laos and Brazil (Simas et al., 2015; Lacroix et al., 2017; Han et al., 2019). Although only partial

* Corresponding author.

E-mail address: LiuGuangliang01@caas.cn (G. Liu).

¹ Jianing Chen and Zemei Wang contributed equally to this work.

genomic information was provided, these findings raise public concern about whether PEDV will be capable of jumping to non-porcine hosts or if it is already in the process of doing so.

In the 21st century, the outbreaks of severe acute respiratory syndrome coronavirus, Middle East respiratory syndrome coronavirus and severe acute respiratory syndrome coronavirus 2 are all related to the cross-species infection of coronaviruses. Therefore, it is important to investigate whether PEDV can infect rodents. In this study, PEDV was detected in rat fecal samples collected from pig farms. BALB/c mice and Sprague Dawley (SD) rats were used for further verification. The results demonstrated that PEDV could cross the species barrier to infect the rodent, providing valuable insights into PEDV behavior.

RESULTS

The PEDV genome is detectable in rat fecal samples collected from pig farms

Infestations associated with rodents have been a persistent challenge for pig farms. Several porcine viruses have previously been isolated in rats (Tonietti et al., 2013; Joshi et al., 2016; Zhai et al., 2016; Murphy et al., 2019; De Sabato et al., 2020). In a prior study, we identified that

the rat crypt epithelial cell line, IEC-6, was highly susceptible to PEDV, suggesting rats may be potentially susceptible to PEDV (Chen et al., 2020). Consequently, we endeavored to further detect PEDV in rat fecal samples collected from pig farms during 2020 and 2022. Among the collected samples, 6 out of 39 were found to be positive for PEDV. Two complete PEDV genomes were then assembled (GenBank access No. OR601542 and No. OR601543; Science Data Bank, No.10.57760/sciencedb.25019). Phenotypic analysis based on the complete genome indicated that the obtained PEDV strains belong to the circulating GIIb cluster (Fig. 1). The information of other reference PEDV strains was listed in Supplementary Table S1. Although PEDV was detected in rat fecal samples, it is challenging to deduce whether rats were infected by PEDV or if the samples were contaminated by PEDV during collection. Therefore, our next step was to attempt to infect mice with PEDV.

Neonatal mice are permissive to PEDV via different inoculation routes

To assess PEDV infection in mice, 7-day-old BALB/c mice were selected and inoculated with PEDV using various routes. Intraperitoneal (i.p.), intragastric (i.g.), oral, and intranasal (i.n.) inoculations were applied with 10^5 TCID₅₀ PEDV per mouse




Fig. 1. Detection and analysis of PEDV genome in rat fecal samples. The complete genome sequences of two PEDV strains were successfully assembled and subjected to phylogenetic analyses. The phylogenetic tree, constructed using the maximum likelihood (ML) method with 1000 bootstrap replicates, is presented. The red arch highlights the assembled PEDV genome from this study.

(Fig. 2A–D). Each group of the survival monitoring experiment contained five mice. In the pathology progression experiment, nine mice were sacrificed at 1, 3, and 5 days post-inoculation (dpi) for viral load determination. The results showed that the i.p. mice and orally inoculated mice displayed no significant clinical symptoms (Fig. 2E and F). One out

of five mice with i.g. died at 5 dpi. The infected mice exhibited anorexia and depression, with watery diarrhea being observed at 2 and 3 dpi in some mice (Fig. 2G). Intranasal mice showed no deaths, but depression and soft feces were observed at 2 and 3 dpi and recovered thereafter (Fig. 2H). All infected mice maintained stable growth during the observation period, except for the i.g. group,




Fig. 2. Neonatal BALB/c mice are susceptible to PEDV infection via various routes. In this study, neonatal BALB/c mice were subjected to PEDV inoculation through different routes: (A) intraperitoneal infection, (B) oral infection, (C) intra gastric infection, and (D) intranasal infection. Following inoculation, we monitored the challenged mice's clinical scores (E, F, G, H), and body weight changes (I, J, K, L) daily for 7 days. Clinical scores were graded as described in [Supplementary Table S3](#). Subsequently, we homogenized the tissues of infected mice for viral load determination. Viral load determination was performed through TCID₅₀ detection (M, N, O, P) and RT-PCR analysis (Q).

which showed significantly slower growth compared to the mock group (Fig. 2I–L).

At 1, 3, and 5 dpi, three mice from each group were euthanized. Different tissues were harvested for viral load determination and RT-PCR analysis. Surprisingly, PEDV was detected among different tissues. Mice in the i.p. (Fig. 2M) had the highest viral titers in the brains and lungs, exceeding 10^2 TCID₅₀/mg tissue. PEDV was also detectable in the lungs, spleens, intestines, and blood for all mice at 3 dpi. In orally inoculated mice, PEDV was only detected in the brains and lungs at 3 dpi (Fig. 2N). Additionally, PEDV was detectable in the brains, lungs, intestines, and blood for i.g. and i.n. mice (Fig. 2O and P). These results were confirmed by RT-PCR analysis. The results revealed strong bands in the brains and lungs, faint bands in the spleens, intestines, or blood, but no signal in other tissues of all inoculated groups at 3 dpi (Fig. 2Q). In summary, all these data suggest that PEDV can infect the brains and lungs of neonatal BALB/c mice via all four different routes.

Neonatal mice are susceptible to PEDV via intracranial inoculation

The preceding results suggest that PEDV can infect neonatal mice at low levels and most efficiently replicate in the brain via i.p., i.g., oral, and i.n. Therefore, our next step was attempting to infect mice with PEDV via intracranial inoculation (i.c.) (Fig. 3A). We infected 7-day-old

mice with PEDV via i.c. (10^5 TCID₅₀ PEDV for each mouse). After inoculation, the PEDV-infected mice exhibited depression and anorexia at 1 dpi (Fig. 3B). Subsequently, it was observed that PEDV infection delayed the growth of infected mice compared to the control group (Fig. 3C). By 5 dpi, 3 out of 5 mice had died, while the remaining mice survived with their body weight increasing from 6 dpi (Fig. 3D). During the observation, the infected mice developed watery diarrhea, and their body weight sharply dropped from 3 dpi (Fig. 3E). Some infected mice even developed opisthotonus. The intestine of infected mice was filled with yellow content and air, contrasting with white chyle-like contents in control mice (Fig. 3F). Although the size of the spleens and the length of the intestines were distinct between infected mice and control mice, there was no significant difference after dividing by body weight (Supplementary Table S2). The results indicated that PEDV was highly lethal to 7-day-old mice via i.c. Additionally, i.c. had nearly little influence on control mice (Supplementary Figs. S1A and 1B). Viral load determination showed that the brain was the most suitable for PEDV growth, with the top viral titers exceeding 10^4 TCID₅₀/mg. PEDV was detectable in the brain, lung, intestine, blood, heart, kidney and feces at 3 and 5 dpi (Fig. 3G). PEDV was also detectable in the spleens at 3 dpi and liver at 5 dpi, with viral titers much higher via i.c. than other inoculation routes. The data above demonstrated that PEDV displays cerebral tropism in neonatal mice.




Fig. 3. PEDV induces lethal outcomes in neonatal BALB/c mice via intracranial inoculation. **A** Seven-day-old BALB/c mice were intracerebrally inoculated with either PEDV or DMEM. The daily observations included **(B)** clinical scores, changes in **(C)** body weight, and **(D)** survival rates. **E** Post-inoculation, PEDV-challenged mice exhibited severe clinical symptoms, particularly watery diarrhea. **F** The intestines of infected mice displayed distension with air and fluidic, containing yellow content, while control mice had undigested, curdled milk masses. **G** The viral load of PEDV in different tissues was determined by TCID₅₀.

More importantly, morbidity and mortality of PEDV in mice vary via different infection routes.

PEDV results in asymptomatic infection in mice older than 7 days

While PEDV can lead to a death rate of over 95% in suckling piglets, it manifests mild symptoms in pigs older than two weeks. To investigate whether PEDV infection in mice is also highly associated with age, 7-day-old, 14-day-old, and 21-day-old mice were used for further examination. The observation period was extended to 14 days for more

detailed monitoring and collection serum for antibody response detection.

During the test, only inactivity, wasting, and depression were observed in 14- and 21-day-old mice (Fig. 4A). The body weight of 14- and 21-day-old mice slightly dropped after inoculation but recovered at 3 dpi (Fig. 4B). No mice older than 7 days died during the observation (Fig. 4C). The 7-day-old mice displayed severe diarrhea symptoms, and three out of five died at 3 dpi. The body weight of the surviving mice differed from that of the control mice. Viral load results revealed that PEDV infection mainly occurred in the brains of 14- and 21-day-old mice




Fig. 4. Age-dependent pathogenicity of PEDV in BALB/c mice. BALB/c mice of varying ages were intracerebrally inoculated with either PEDV or DMEM. Daily observations included (A) clinical scores, (B) changes in body weight, and (C) survival rates. Post-inoculation, PEDV-challenged mice were euthanized, and tissues were harvested at different days post-inoculation (dpi). The viral load of PEDV in different tissues was determined by TCID₅₀ for (D) 7-day-old, (E) 14-day-old, and (F) 21-day-old mice. Additionally, serum samples were collected from (G) 7-day-old, (H) 14-day-old, and (I) 21-day-old mice for detecting PEDV-specific IgG at 14 dpi.

but with lower viral loads compared to 7-day-old mice (Fig. 4D–F). PEDV was not detected in other organs of older mice, except in the lung of one 21-day-old mouse at 5 dpi. The viral titers in the brain also peaked at 3 dpi for all infected mice. At 14 dpi, PEDV was not detectable in the brains of 21-day-old mice. Additionally, seroconversion was detected in 7-day-old mice but not in older mice (Fig. 4G–I). In conclusion, PEDV also displayed age-related morbidity and mortality in mice via i.c.

The mortality of PEDV in mice is viral dose-dependent

To determine the lowest dose that can result in death, 7-day-old mice were inoculated with PEDV at different doses (10^5 , 10^4 , 10^3 , 10^2 , and 10^1 TCID₅₀) or DMEM. Each group contained three co-housed mice, which were also sacrificed at 14 dpi to assess the transmission of PEDV.

The results demonstrated that mice inoculated with 10^5 , 10^4 , and 10^3 TCID₅₀ PEDV displayed severe watery diarrhea (Fig. 5A). The death rates for these three groups were 60%, 40%, and 40%, respectively (Fig. 5B). All deaths occurred before 5 dpi. Although mice inoculated with 10^2 and 10^1 TCID₅₀ viruses appeared asymptomatic, one mouse inoculated with 10^2 TCID₅₀ viruses died at 11 dpi (Fig. 5A and B). The size and body weight of the mice inoculated with 10^2 or 10^1 TCID₅₀ viruses were larger than that of three other groups (Fig. 5C and D).

The viral titration results showed that the replication of PEDV peaked at 3 dpi and decreased thereafter for all groups. At 14 dpi, PEDV could be detected in the brains, lungs, intestines, blood, and feces of mice inoculated with more than 10^3 TCID₅₀, suggesting a potential source of infection (Fig. 5E and F). For mice inoculated with 10^3 or 10^2 TCID₅₀ viruses, PEDV could be detected in the brains, lungs, intestines, and spleens before 7 dpi and occasionally in other tissues (Fig. 5G and H). For mice inoculated with 10^1 TCID₅₀ PEDV, the virus could only be detected in the brains and lungs before 7 dpi (Fig. 5I).

Seroconversion was then confirmed in inoculated mice, with OD₄₅₀ nm values around 0.4–0.7, compared to about 0.2 for control mice (Fig. 5J). Although no virus was detected in co-housed mice at 14 dpi, antibodies against PEDV-nucleocapsid protein were detectable in co-housed mice, except for mice co-housed with 10^1 and 10^2 TCID₅₀ inoculated mice (Fig. 5K) but not in the parent mice (Fig. 5L).

PEDV of different genotypes are fatal to neonatal mice

PEDV has been categorized into two lineages, distinguished by differences in immunity and pathogenicity for piglets. The classic lineage, also known as genotype I, was initially identified in the 1970s and rarely detected since the 2010s, with the representative strain being CV777. The genotype II virus has been prevalent worldwide since 2013, and in the above studies, the genotype II LJX strain was used for examination. Here, the genotype I CV777 strain was also utilized to test for any differences.

7-day-old BALB/c mice were intracranially inoculated with 10^5 TCID₅₀ per mouse. The inoculated mice developed severe watery diarrhea at 1 dpi, and 3 out of 5 mice exhibited opisthotonus (Fig. 6A and B). Their body weight did not increase after inoculation (Fig. 6C), and all mice died within 5 dpi (Fig. 6D). Both the small and large intestines were filled with yellow content and air, while the large intestine turned transparent (Fig. 6E). Viral loading determination demonstrated the presence of the virus in various tissues at 3 and 5 dpi (Fig. 6F). The combination of high mortality and high viral loading in tissues suggests that PEDV from different genotypes is also destructive to neonatal mice.

PEDV infection induces pathological changes in neonatal mice

The histopathological alterations in various tissues associated with PEDV infection were then determined in mice. The results revealed that PEDV infection in the brain led to pronounced neuronal edema. Furthermore, a satellite phenomenon was observed, wherein neurons were surrounded by oligodendrocytes, suggesting neuronal apoptosis.

Additionally, the formation of cerebral thrombosis was significantly higher in infected mice (Fig. 7A). Control mice, on the other hand, showed no obvious lesions, implying no damage caused by intracranial inoculation. In the lungs of infected mice, serious interstitial pneumonia was observed, characterized by thickening of the alveolar wall and achrocytosis. These findings suggest that PEDV also induces severe respiratory symptoms. Although the virus was detected in the intestine and spleen, only mild lesions were observed, with inclusion bodies accumulating primarily in the spleen (Fig. 7A). The immunohistochemistry analysis was performed to detect viral distribution using antibodies against the structural protein N or nonstructural protein 16 (Fig. 7B). PEDV was found to spread throughout the brain, with a high concentration in the cerebral cortex and fewer signals in the fornix. In the lung, strong signals were observed in the trachea and alveolar wall, indicating PEDV infection in the respiratory tract. Moreover, PEDV infected the intestine and spleen, with signals detected in the intestinal villus and medulla of the spleen. In conclusion, PEDV can result in systemic infection in mice by intracranial inoculation, causing severe lesions in the brain and lung but milder effects in other tissues.

PEDV results in asymptomatic infection in sucking rats

As the mouse is identified to be susceptible to PEDV, we then test whether PEDV can infect the rat. Similar, 7-day-old rats were inoculated with 10^5 TCID₅₀ PEDV via i.p., i.g., oral, i.n. and i.c. All infected rats were alive and displayed no observable clinical symptoms during the observation for 7 days, which was distinct to mice. The viral titration assays showed that PEDV was also detectable among the brains, lungs, and intestines for each inoculation route before 5 dpi (Fig. 8A–E). PEDV was also detectable within the spleens, intestines and feces of rats derived from oral, i.g. and i.c. group at 7 dpi. The viral titers in rat intestines were higher than that in mouse intestines, suggesting rats may be susceptible to natural infection of PEDV via fecal-oral transmission than mice. Immunohistochemistry analysis also supported the results above. The positive signals were observed within the brains, lungs, intestines, and spleen. Although the signals within the rat brains and lungs were significantly reduced comparing to mouse, they were enhanced in the intestines and spleens, potentially explaining why PEDV was not lethal to rats (Fig. 8F).

PEDV infection is confirmed in the brains of piglets

While this study has identified that PEDV targets the brains of the rodent, this manifestation has not been verified in pigs. To gain a deeper understanding of PEDV infection, we selected three 5-day-old piglets and infected them via oral inoculation with the PEDV LJX strain. All pigs developed severe clinical symptoms and succumbed to the infection by 4 dpi. The real-time PCR analysis revealed over 10^8 copies/mg of the PEDV genome in the intestine, whereas a relatively low number of PEDV genome copies were detected in pig brains and other tissues (Fig. 9A). The PEDV genome copies within the brains and other tissues closely approached the cut-off line. However, immunohistochemical staining demonstrated the presence of PEDV in the brain, identified among the neurons, leading to their necrosis in all three piglets, while positive signals were not observed in mock-infected pigs (Fig. 9B). Finally, Vero cells were infected with the filtered brain homogenate and analyzed with immunofluorescence assay (IFA). The presence of positive signals demonstrated the existence of infectious PEDV virions within the pig brain (Fig. 9C).

DISCUSSION

PEDV is an enteric virus belonging to the genus *Alphacoronavirus* within the *Coronaviridae* family. This virus infects pigs of all ages, causing mortality rates of up to 100% in suckling piglets. Since its first identification in the 1970s, PEDV has spread worldwide. Previous studies have established that pigs are the only identified natural reservoir for PEDV.




Fig. 5. Dose-related morbidity and mortality in BALB/c mice upon intracerebral inoculation. 7-day-old BALB/c mice were intracerebrally inoculated with varying doses of PEDV or DMEM. Daily observations included (A) clinical scores, (B) survival rates, and (C) changes in body weight. D At 14 dpi, the size and condition of surviving mice varied among different groups. Post-inoculation, PEDV-challenged mice were euthanized, and tissues were harvested at 1, 3, 5, 7, and 14 dpi. The viral load of PEDV in different tissues for mice inoculated with (E) 10⁵ TCID₅₀, (F) 10⁴ TCID₅₀, (G) 10³ TCID₅₀, (H) 10² TCID₅₀, and (I) 10¹ TCID₅₀ was determined. Additionally, serum samples were collected from (J) PEDV-infected mice, (K) co-housed mice and (L) parent mice for detecting PEDV-specific IgG at 14 dpi.




Fig. 6. Genotype I PEDV exhibits enhanced-pathogenicity in neonatal mice. Seven-day-old BALB/c mice were intracerebrally inoculated with PEDV CV777 strain or DMEM. A Post-inoculation, CV777-challenged mice exhibited severe clinical symptoms from 1 dpi. Daily observations included (B) clinical scores, (C) changes in body weight, and (D) survival rates. E The small intestine of infected mice showed signs of air and fluid distension with a transparent appearance in the large intestine. F Viral titers of CV777 in different tissues were determined through TCID₅₀ analysis.

However, in this study, we detected the PEDV genome in rat fecal samples and further discovered that mice and rats could be infected by PEDV via different routes. With variations in inoculation routes and doses, PEDV can result in either asymptomatic or fatal infections. Notably, PEDV infection in rodents differs from its manifestation in piglets, targeting the brains, lungs, and spleens. These findings indicate that rodents may be susceptible to PEDV.

Most coronaviruses exhibit a limited host range. However, those with the capacity for cross-species transmission can lead to disasters in both humans and animals. Therefore, it is crucial to investigate the transmission of coronaviruses among different species. Mice are commonly used as an animal model for pathogenic and transmission evaluations. Several coronaviruses, traditionally with limited hosts, have been proven to replicate in mice, including HCoV-OC43, swine acute diarrhea syndrome coronavirus (SADS-CoV), porcine hemagglutinating encephalomyelitis virus (PHEV), and even the avian coronavirus infectious bronchitis virus (IBV) (McIntosh et al., 1967; Butler et al., 2006; Mora-Díaz et al., 2021; Chen et al., 2022). However, the clinical symptoms observed in mice differ significantly from those in the natural reservoirs, often involving severe neurological signs and high viral titers in the

central nervous system. These studies suggest that coronaviruses may change when encountering new hosts, evading detection.

Neonatal mice have been experimentally infected with PEDV via intracranial inoculation in previous studies, yielding low morbidity and mortality in mice aged three days or younger (Kotani et al., 2013). However, the details of the infection were poorly characterized, with PEDV only detectable in the brain and not in other tissues. Moreover, transgenic mouse expressing porcine aminopeptidase N was also developed and proved to be susceptible to PEDV (Park et al., 2015). This study shows that PEDV can infect both mice and rats through various routes, leading to a range of manifestations from asymptomatic to fatal infections.

Our study also identified PEDV infection in pig brains in the comparative test. The viral titers were relatively low in the brain compared to the intestine, but infectious virions were present, causing neuronal necrosis. This manifestation has not been observed before and may potentially explain why the production capacity of PEDV-infected sows is unstable. It is also intriguing to note another swine coronavirus, PHEV, which primarily causes neurological signs but occasionally diarrhea for some variants (Mora-Díaz et al., 2019). Could PEDV




Fig. 7. Pathogenic analysis of PEDV-infected mice. Seven-day-old BALB/c mice were intracerebrally inoculated with PEDV or DMEM and sacrificed at 3 dpi. The brain, lung, intestine, and spleen samples were collected and subjected to (A) H&E staining or (B) immunohistochemistry. Edema in the brain and lymphocyte infiltration in the lung were indicated by yellow arrows. The satellite phenomenon observed in the brain and the thickening of interstitial tissue in the lung were highlighted by red arrows. Thrombus formations were marked by black arrows, and inclusion bodies in the spleen were pointed out by red arrows. PEDV particles in different tissues were identified by red arrows in the immunohistochemistry images.




Fig. 8. PEDV infection in 7-day-old rats. 7-day-old Sprague Dawley (SD) rats were inoculated with PEDV or DMEM via different routes for survival test or viral titer determination: (A) intraperitoneal infection, (B) intra gastric infection, (C) oral infection, (D) intranasal infection and (E) intracranial infection. F Although no obvious lesions were observed, immunohistochemistry staining showed the presence of PEDV particles in different tissues, which were indicated by red arrows.




Fig. 9. Identification of PEDV infection in the brains of piglets. Three 5-day-old piglets were orally inoculated with PEDV, resulting in mortality by 4 dpi. One piglet was inoculated with an equal volume of DMEM as a control. **A** Various tissues, including the brain, were harvested for viral load determination using real-time PCR analysis. **B** Additionally, the brains were subjected to immunohistochemical staining to identify positive signals (red arrow). **C** Vero cells were inoculated with the filtered brain homogenate and analyzed using immunofluorescence assay (IFA) to confirm the presence of infectious PEDV virions.

occasionally switch its characteristics in some variants? PEDV variants, with a large deletion among the spikes, could infect both the lungs and intestines, which were observed previously. Similar mutations and results were also observed in transmissible gastroenteritis virus (TGEV). A large deletion within the TGEV spike also switched its tissue tropism to infect and replicate in the lung. It is predictable that the PEDV variants will bring more uncertainty in the future.

Several swine viruses have been identified and isolated in rodents on pig farms. Porcine circovirus type 2 (PCV2), an economically important swine pathogen, has been reported to frequently spillover from pigs to rats on pig farms (Lorincz et al., 2010; Pinheiro et al., 2013; Zhai et al., 2016). Senecavirus A (SVA), an emerging picornavirus with swine as natural reservoirs, has also been detected and isolated from rat feces and small intestines collected from pig farms in the USA (Joshi et al., 2016). Additionally, rats captured from pig farms carrying the hepatitis E virus (HEV) have drawn increasing attention (Sanford et al., 2012; De Sabato et al., 2020; Wang and Meng, 2021). Although pigs and wild boars were previously considered the only natural reservoirs for PCV2, the high positive

rate of HEV in rats on pig farms complicates its control and prevention. Rodents seem to be a potential source for spilling porcine viruses to other species. Importantly, these viruses cause serious clinical symptoms in pigs but are often asymptomatic in rats. Therefore, their infection in rats is frequently overlooked, and the pathogenicity and transmission of these viruses in rats are poorly defined. Similar to these viruses, our study identified that PEDV infection turned out to be asymptomatic to rats. Although the replication was observed, no rats died via different infection routes, suggesting why PEDV may be ignored in rats.

The differences in immune responses and physiological conditions between the rodents and pigs might influence the phenotypes upon PEDV infection. Both sucking mice and rats have undeveloped blood-brain barriers, which may facilitate the entry and spread of PEDV. Pigs have more complex barriers between different tissues. Additionally, sucking pigs have better developed immune systems when compared to the rodents, which may further restrict the growth and spread of PEDV. These differences will primarily contribute to the distinct infection results between the rodents and pigs.

CONCLUSIONS

To our knowledge, this study provides the first evidence of PEDV detection in rat fecal samples and robust data supporting its cross-species transmission potential. Although viral isolation from field-collected rat feces and capture of naturally infected rats were unsuccessful—approaches that would better reflect natural transmission dynamics—we successfully established neonatal rodents as viable models. We further found that PEDV resulted in systemic infection in mice and rats, with morbidity and mortality rates demonstrating significant dose- and route-dependence. While efficient transmission was not observed, our findings confirm that PEDV have acquired the capacity to replicate in rodent. Continuous surveillance of PEDV evolution is therefore warranted to mitigate zoonotic risks.

MATERIALS AND METHODS

Virus and cells

Vero-E6 cells were purchased from the China National collection of authenticated cell cultures and maintained in our lab. The cells were cultured with DMEM (Sigma, Germany) supplemented with 10% fetal bovine serum (Sigma, Germany) and 100 IU of penicillin per mL at 37 °C in the presence of 5% CO₂. The PEDV LJX01/GS/2014 and CV777 strains used in this study were propagated in our laboratory. LJX01/GS/2014, belonging to gene type II, shared nucleotide homology of 99.9% and 97.9% with the identified two strains. The CV777 strain was a representative strain of gene type I viruses, significantly different from the identified strains.

Sample collection and detection

The rat fecal samples were collected from family pig farms between 2020 and 2022. All samples were collected from different areas of the family pig farms but away from the pig house to reduce the potential contaminants. Only fresh rat fecal sample free of visible contaminants were collected. After collection, the samples were dissolved in sterile PBS and then detected by RT-PCR analysis.

Rodent infection experiments

The BALB/c mice and SD rats of 7-day-old were purchased from the Experimental Animal Research Center of Lanzhou Veterinary Research Institute (LVRI). Animal experiments followed the recommendations of the Guide for the Care and Use of Laboratory Animals of LVRI. All rodents used in this study were detected to be free of PEDV, mouse hepatitis virus and murine rotavirus and kept in the Experimental Animal Research Center of LVRI, which could ensure the absence of exogenous pathogens. The experiments included two independent sections: a survival monitoring experiment and a pathology progression experiment. The former included five mice or rats for each group. The latter included fifteen mice each group for a 7-day test and twenty mice each group for a 14-day test. And 10 rats for each group were used in the viral titer determination. To evaluate PEDV infection, rodents were inoculated via intraperitoneal (i.p.), intragastric administration (i.g.), oral, intranasal (i.n.), or intracranial (i.c.) routes with a 50 µL volume for each one. The control animals were inoculated with an equal volume of DMEM. All animals were observed daily for clinical signs, and survival. The severity of clinical symptoms was scored on a scale of five grades (see [Supplementary Table S3](#)).

Pig infection experiment

Four 5-day-old Landrace piglets, detecting to be free of PEDV, transmissible gastroenteritis virus (TGEV), porcine deltacoronavirus (PDCoV), African swine fever virus (ASFV) and porcine reproductive and

respiratory syndrome virus (PRRSV), were purchased from a pig farm and kept in the Experimental Animal Research Center of LVRI, which could ensure the absence of exogenous pathogens. Three pigs were orally inoculated with 10⁶ TCID₅₀ PEDV LJX strain, while one served as negative control, inoculated with an equal volume of DMEM. Infected piglets died at 4 dpi, and the control was then euthanized. All tissues were harvested for viral loading determination. Brains were also prepared for immunohistochemical staining.

Viral load determination

For viral load determination, rodents from each group were euthanized by CO₂ at 1, 3, 5, 7, and 14 dpi. The blood was collected by cutting tails of the tested mice before euthanasia. The tissues, were weighed, and mixed with 1 mL PBS in ice waiting for homogenation within 15 min. The homogenates were then centrifuged at 12 000 ×g for 10 min at 4 °C, and filtered with 0.45 µm filters. The filtrate was serially diluted (10⁻¹ to 10⁻⁶) and inoculated in Vero-E6 cells. After 1 h post-infection, cells were washed three times with PBS and incubated for a further 72 h. Viral titers were checked by immunofluorescence assay (IFA) with a fluorescence microscope (TE2000U; Nikon) and calculated by the Reed-Muench method.

Immunofluorescence assay

Vero-E6 cells were seeded in 96-well plates. At 85% confluency, cells were infected with filtered tissue homogenate. After 48 h, cells were treated with 4% paraformaldehyde for fixing, 0.5% Triton X-100 for permeation, and 5% skim milk for blocking. Incubation with mouse anti-PEDV-nucleocapsid monoclonal antibody (1:5000 dilution) and DyLight 488 goat anti-mouse IgG antibody (1:1000 dilution) (Biodragon, China) followed. Cells were examined under a fluorescence microscope (TE2000U; Nikon) with a video documentation system.

Antibody response against PEDV infection

Serum collected from PEDV-infected mice, co-housed mice, parent mice, and control mice at 14 dpi was used for ELISA. The serum (1:50 dilution) was incubated with an ELISA plate coated with PEDV-nucleocapsid protein (100 ng per well) at 37 °C for 1 h. After three washes with PBS, HRP-conjugated goat anti-mouse IgG (H + L) (1:500 dilution) was added, and absorbance values were scanned at OD₄₅₀ nm with a microplate reader.

RT-PCR and real-time PCR analysis

RT-PCR analysis for PEDV detection used 2× TsingKe Master Mix (TsingKe, China). Reactions were incubated at 95 °C for 3 min and subjected to 35 cycles of 95 °C for 15 s, 52 °C for 15 s, and 72 °C for 30 s, with a final extension step of 72 °C for 5 min. PCR products were analyzed by 1% agarose gel electrophoresis. Real-time qPCR for PEDV genome copies detection used Universal U + Probe Master Mix V2 (Vazyme, China) by the Bio-Rad CFX96 system, which was described in previous study ([Huang et al., 2019](#)). Reactions were incubated at 94 °C for 30 s, followed by 45 cycles at 94 °C for 5 s and 60 °C for 30 s.

Histopathology and immunohistochemical staining

The rodents infected with PEDV or uninfected were euthanized by CO₂ at 3 dpi for histopathologic and immunohistochemical examination. Tissues were immediately harvested, fixed in 10% formalin at room temperature for at least 48 h, bisected, and embedded in paraffin. Sections were deparaffinized, rehydrated by xylene three times, and dehydrated by ethanol of different concentrations. For histopathologic examination, tissue sections were stained with hematoxylin and eosin. For immunohistochemical testing, tissue sections were dewaxed and

dehydrated. After deparaffination, sections were treated with 3% H₂O₂ for 10 min, followed by distilled water washing and microwave treatment in a citrate buffer for 20 min at 99 °C. Sections were then blocked with 5% BSA for 30 min and further incubated with mouse anti-PEDV-nucleocapsid antibodies (1:1000 dilution) or rabbit anti-PEDV-nsp16 antibodies (1:500 dilution) for 1 h at 37 °C, followed by biotinylated goat anti-mouse antibodies (1:500 dilution) or goat anti-rabbit antibodies (1:500 dilution) (Abbkine, China) for 30 min at room temperature. Sections were treated by 3, 3'-diaminobenzidine tetrahydrochloride chromogen (Beyotime, China), counterstained with hematoxylin and visualized by a light microscope.

Statistical analysis

For statistical significance examination, the Student's *t*-test was used between matched groups. An unadjusted *P* value of less than 0.05 was considered significant, and a *P* value of less than 0.01 was considered extremely significant.

DATA AVAILABILITY

All the data generated during the current study are included in the manuscript.

ETHICS STATEMENT

The animal experiment was conducted in compliance with the Animal Ethics Procedures and Guidelines of Laboratory Animals, approved by the Animal Administration and Ethics Committee of Lanzhou Veterinary Research Institute (LVRI) of the Chinese Academy of Agricultural Sciences (Permit No. LVRIAEC-2022-076).

AUTHOR CONTRIBUTIONS

Jianing Chen: data curation, writing-original draft, project administration. Zemei Wang, Shengyu Lin, Yongheng Shao, Mengling Gao, Shuxian Li, Qingbo Chen, Yonghao Hu: data curation. Guangliang Liu: project administration, writing-review & editing.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

ACKNOWLEDGEMENTS

This work was supported by the Basic Research Innovation Group Project of Gansu Province (No. 25JRRA434) and the National Natural Science Foundation of China (U22A20522).

APPENDIX A. SUPPLEMENTARY DATA

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.virs.2025.03.012>.

REFERENCES

- Butler, N., Pewe, L., Trandem, K., Perlman, S., 2006. Murine encephalitis caused by HCoV-OC43, a human coronavirus with broad species specificity, is partly immune-mediated. *Virology* 347, 410–421.
- Chen, J., Cui, Y., Wang, Z., Liu, G., 2020. Identification and characterization of PEDV infection in rat crypt epithelial cells. *Vet. Microbiol.* 249, 108848.
- Chen, Y., Jiang, R.D., Wang, Q., Luo, Y., Liu, M.Q., Zhu, Y., Liu, X., He, Y.T., Zhou, P., Yang, X.L., Shi, Z.L., 2022. Lethal swine acute diarrhea syndrome coronavirus infection in suckling mice. *J. Virol.* 96, e0006522.
- Chung, H.C., Lee, J.H., Nguyen, V.G., Huynh, T.M.L., Lee, G.E., Moon, H.J., Park, S.J., Kim, H.K., Park, B.K., 2016. New emergence pattern with variant porcine epidemic diarrhea viruses, South Korea, 2012–2015. *Virus Res.* 226, 14–19.
- De Sabato, L., Ianiro, G., Monini, M., De Lucia, A., Ostanello, F., Di Bartolo, I., 2020. Detection of hepatitis E virus RNA in rats caught in pig farms from Northern Italy. *Zoonoses Public Health* 67, 62–69.
- Han, Y., Du, J., Su, H., Zhang, J., Zhu, G., Zhang, S., Wu, Z., Jin, Q., 2019. Identification of diverse bat alphacoronaviruses and betacoronaviruses in China Provides new insights into the evolution and origin of coronavirus-related diseases. *Front. Microbiol.* 10, 1900.
- Huang, X., Chen, J., Yao, G., Guo, Q., Wang, J., Liu, G., 2019. A TaqMan-probe-based multiplex real-time RT-qPCR for simultaneous detection of porcine enteric coronaviruses. *Appl. Microbiol. Biotechnol.* 103, 4943–4952.
- Joshi, L.R., Mohr, K.A., Clement, T., Hain, K.S., Myers, B., Yaros, J., Nelson, E.A., Christopher-Hennings, J., Gava, D., Schaefer, R., et al., 2016. Detection of the emerging Picornavirus Senecavirus A in pigs, mice, and houseflies. *J. Clin. Microbiol.* 54, 1536–1545.
- Kotani, O., Shirato, K., Nagata, N., Ikeda, H., Takahashi, K., Taguchi, F., 2013. Neuropathogenesis of a mouse-adapted porcine epidemic diarrhea virus infection in suckling mice. *J. Gen. Virol.* 94, 831–836.
- Lacroix, A., Duong, V., Hul, V., San, S., Davun, H., Omaliss, K., Chea, S., Hassanin, A., Theppangna, W., Silithammavong, S., et al., 2017. Genetic diversity of coronaviruses in bats in Lao PDR and Cambodia. *Infect. Genet. Evol.* 48, 10–18.
- Lorincz, M., Cságha, A., Biksi, I., Szeredi, L., Dán, A., Tuboly, T., 2010. Detection of porcine circovirus in rodents - short communication. *Acta Vet. Hung.* 58, 265–268.
- Lv, L., Luo, H., Yu, L., Tong, W., Jiang, Y., Li, G., Tong, G., Li, Y., Liu, C., 2022. Identification and characterization of cell lines HepG2, Hep3B217 and SNU387 as models for porcine epidemic diarrhea coronavirus infection. *Viruses* 14, 2754.
- McIntosh, K., Becker, W.B., Chanock, R.M., 1967. Growth in suckling-mouse brain of "TBV-like" viruses from patients with upper respiratory tract disease. *Proc. Natl. Acad. Sci. U. S. A.* 58, 2268–2273.
- Mora-Díaz, J.C., Piñeyro, P.E., Houston, E., Zimmerman, J., Giménez-Lirola, L.G., 2019. Porcine hemagglutinating encephalomyelitis virus: a review. *Front. Vet. Sci.* 6, 53.
- Mora-Díaz, J.C., Piñeyro, P.E., Rauh, R., Nelson, W., Sankoh, Z., Gregg, E., Carrillo-Ávila, J.A., Shen, H., Nelli, R.K., Zimmerman, J.J., et al., 2021. Porcine hemagglutinating encephalomyelitis virus infection *in vivo* and *ex vivo*. *J. Virol.* 95, e02335-20.
- Murphy, E.G., Williams, N.J., Bennett, M., Jennings, D., Chantrey, J., McElhinney, L.M., 2019. Detection of Seoul virus in wild brown rats (*Rattus norvegicus*) from pig farms in northern England. *Vet. Rec.* 184, 525.
- Niu, Z., Zhang, S., Xu, S., Wang, J., Wang, S., Hu, X., Zhang, L., Ren, L., Zhang, J., Liu, X., et al., 2023. Porcine epidemic diarrhea virus replication in human intestinal cells reveals potential susceptibility to cross-species infection. *Viruses* 15, 956.
- Park, J.E., Park, E.S., Yu, J.E., Rho, J., Paudel, S., Hyun, B.H., Yang, D.K., Shin, H.J., 2015. Development of transgenic mouse model expressing porcine aminopeptidase N and its susceptibility to porcine epidemic diarrhea virus. *Virus Res.* 197, 108–115.
- Pensaert, M.B., de Bouck, P., 1978. A new coronavirus-like particle associated with diarrhea in swine. *Arch. Virol.* 58, 243–247.
- Pinheiro, A.L., Bulos, L.H., Onofre, T.S., de Paula, GabardoM., de Carvalho, O.V., Fausto, M.C., Guedes, R.M., de Almeida, M.R., Silva, JúniorA., 2013. Verification of natural infection of peridomestic rodents by PCV2 on commercial swine farms. *Res. Vet. Sci.* 94, 764–768.
- Sanford, B.J., Opriessnig, T., Kenney, S.P., Dryman, B.A., Córdoba, L., Meng, X.J., 2012. Assessment of the cross-protective capability of recombinant capsid proteins derived from pig, rat, and avian hepatitis E viruses (HEV) against challenge with a genotype 3 HEV in pigs. *Vaccine* 30, 6249–6255.
- Simas, P.V., Barnabé, A.C., Durães-Carvalho, R., Neto, D.F., Caserta, L.C., Artacho, L., Jacomassa, F.A., Martini, M.C., Bianchi, Dos, Santos, M.M., et al., 2015. Bat coronavirus in Brazil related to appalachian ridge and porcine epidemic diarrhea viruses. *Emerg. Infect. Dis.* 21, 729–731.
- Stevenson, G.W., Hoang, H., Schwartz, K.J., Burrough, E.R., Sun, D., Madson, D., Cooper, V.L., Pillatzki, A., Gauger, P., Schmit, B.J., et al., 2013. Emergence of Porcine epidemic diarrhea virus in the United States: clinical signs, lesions, and viral genomic sequences. *J. Vet. Diagn. Invest.* 25, 649–654.
- Sun, R., Cai, R., Chen, Y., Liang, P., Chen, D., Song, C., 2012. Outbreak of porcine epidemic diarrhea in suckling piglets, China. *Emerg. Infect. Dis.* 18, 161–163.
- Sun, Y., Chen, Y., Han, X., Yu, Z., Wei, Y., Zhang, G., 2019. Porcine epidemic diarrhea virus in Asia: an alarming threat to the global pig industry. *Infect. Genet. Evol.* 70, 24–26.
- Tonietti, Pde, O., da Hora, A.S., Silva, F.D., Ferrari, K.L., Brandão, P.E., Richtzenhain, L.J., Gregori, F., 2013. Simultaneous detection of group A rotavirus in swine and rat on a pig farm in Brazil. *Sci. World J.* 2013, 648406.
- Wang, B., Meng, X.J., 2021. Hepatitis E virus: host tropism and zoonotic infection. *Curr. Opin. Microbiol.* 59, 8–15.
- Wang, X., Fang, L., Liu, S., Ke, W., Wang, D., Peng, G., Xiao, S., 2019. Susceptibility of porcine IPI-2I intestinal epithelial cells to infection with swine enteric coronaviruses. *Vet. Microbiol.* 233, 21–27.
- Wood, E.N., 1977. An apparently new syndrome of porcine epidemic diarrhoea. *Vet. Rec.* 100, 243–244.
- Zhai, S., Chen, S., Liu, W., Li, X., Deng, S., Wen, X., Luo, M., Lv, D., Wei, W., Chen, R., 2016. Molecular detection and genome characterization of porcine circovirus type 2 in rats captured on commercial swine farms. *Arch. Virol.* 161, 3237–3244.
- Zhang, J., Guo, L., Xu, Y., Yang, L., Shi, H., Feng, L., Wang, Y., 2017. Characterization of porcine epidemic diarrhea virus infectivity in human embryonic kidney cells. *Arch. Virol.* 162, 2415–2419.
- Zhao, S., Gao, J., Zhu, L., Yang, Q., 2014. Transmissible gastroenteritis virus and porcine epidemic diarrhoea virus infection induces dramatic changes in the tight junctions and microfilaments of polarized IPEC-J2 cells. *Virus Res.* 192, 34–45.